

# Improved Bounds for Wireless Localization

Tobias Christ<sup>1</sup>, Michael Hoffmann<sup>1</sup>, Yoshio Okamoto<sup>2</sup>, and Takeaki Uno<sup>3</sup>

<sup>1</sup> Institute for Theoretical Computer Science, ETH Zürich, Switzerland

<sup>2</sup> Tokyo Institute of Technology, Japan

<sup>3</sup> National Institute of Informatics, Tokyo, Japan

**Abstract.** We consider a novel class of art gallery problems inspired by wireless localization. Given a simple polygon  $P$ , place and orient guards each of which broadcasts a unique key within a fixed angular range. Broadcasts are not blocked by the edges of  $P$ . The interior of the polygon must be described by a monotone Boolean formula composed from the keys. We improve both upper and lower bounds for the general setting by showing that the maximum number of guards to describe any simple polygon on  $n$  vertices is between roughly  $\frac{3}{5}n$  and  $\frac{4}{5}n$ . For the natural setting where guards may be placed aligned to one edge or two consecutive edges of  $P$  only, we prove that  $n - 2$  guards are always sufficient and sometimes necessary.


## 1 Introduction

Art gallery problems are a classic topic in discrete and computational geometry, dating back to the question posed by Victor Klee in 1973: “How many guards are necessary, and how many are sufficient to patrol the paintings and works of art in an art gallery with  $n$  walls?” Chvátal [2] was the first to show that  $\lfloor n/3 \rfloor$  guards are always sufficient and sometimes necessary, while the beautiful proof of Fisk [6] made it into “the book” [1]. Nowadays there is a vast literature [12, 14, 16] about variations of this problem, ranging from optimization questions (minimizing the number of guards [10] or maximizing the guarded boundary [7]) over special types of guards (mobile guards [11] or vertex pi-guards [15]) to special types of galleries (orthogonal polygons [8] or curvilinear polygons [9]).

A completely different direction has recently been introduced by Eppstein, Goodrich, and Sitchinava [5]. They propose to modify the concept of visibility by not considering the edges of the polygon/gallery as blocking. The motivation for this model stems from communication in wireless networks where the signals are not blocked by walls, either. For illustration, suppose you run a café (modeled, say, as a simple polygon  $P$ ) and you want to provide wireless Internet access to your customers. But you do not want the whole neighborhood to use your infrastructure. Instead, Internet access should be limited to those people who are located within the café. To achieve this, you can install a certain number of devices, let us call them guards, each of which broadcasts a unique (secret) key in an arbitrary but fixed angular range. The goal is to place guards and adjust their angles in such a way that everybody who is inside the café can

prove this fact just by naming the keys received and nobody who is outside the café can provide such a proof. Formally this means that  $P$  can be described by a monotone Boolean formula over the keys, that is, a formula using the operators AND and OR only, negation is not allowed. It is convenient to model a guard as a subset of the plane, namely the area where the broadcast from this guard can be received. This area can be described as an intersection or union of at most two halfplanes. Using this notation, the polygon  $P$  is to be described by a combination of the operations union and intersection over the guards. For example, the first polygon to the right can be described by  $(a \cup b) \cap c \cap d$ .

**Natural guards.** Natural locations for guards are the vertices and edges of the polygon. A guard which is placed at a vertex of  $P$  is called a *vertex guard*. A vertex guard is *natural* if it covers exactly the interior angle of its vertex. But natural vertex guards alone do not always suffice [5], as the second polygon  $P$  shown to the right illustrates: No natural vertex guard can distinguish the point  $p$  inside  $P$  from the point  $q$  outside of  $P$ . A guard placed anywhere on the line given by an edge of  $P$  and broadcasting within an angle of  $\pi$  to the inner side of the edge is called a *natural edge guard*. Dobkin, Guibas, Hershberger, and Snoeyink [4] showed that  $n$  natural edge guards are sufficient for any simple polygon with  $n$  edges.



**Vertex guards.** Eppstein et al. [5] proved that any simple polygon with  $n$  edges can be guarded using at most  $n - 2$  (general, that is, not necessarily natural) vertex guards. More generally, they show that  $n + 2(h - 1)$  vertex guards are sufficient for any simple polygon with  $n$  edges and  $h$  holes. This bound is not known to be tight. Damian, Flatland, O'Rourke, and Ramaswami [3] describe simple polygons with  $n$  edges which require at least  $\lfloor 2n/3 \rfloor - 1$  vertex guards.

**General guards.** In the most general setting, we do not have any restriction on the placement and the angles of guards. So far the best upper bound known has been the same as for vertex guards, that is,  $n - 2$ . On the other hand, if the polygon does not have collinear edges then at least  $\lceil n/2 \rceil$  guards are always necessary [5]. The lower bound construction of Damian et al. [3] for vertex guards does not provide an improvement in the general case, where these polygons can be guarded using at most  $\lceil n/2 \rceil + 1$  guards. As O'Rourke wrote [13]: "The considerable gap between the  $\lceil n/2 \rceil$  and  $n - 2$  bounds remains to be closed."

**Results.** We provide a significant step in bringing the two bounds for general guards closer together by improving both on the upper and on the lower side. On one hand we show that for any simple polygon with  $n$  edges  $\lfloor (4n - 2)/5 \rfloor$  guards are sufficient. The result generalizes to polygons combined in some way by the operations intersection and/or union. Any simple polygon with  $h$  holes can be guarded using at most  $\lfloor (4n - 2h - 2)/5 \rfloor$  guards. On the other hand we describe a family of polygons which require at least  $\lceil (3n - 4)/5 \rceil$  guards. Furthermore, we extend the result of Dobkin et al. [4] to show that  $n - 2$  natural (vertex or edge) guards are always sufficient. It turns out that this bound is tight.

**Table 1.** Number of guards needed for a simple polygon on  $n$  vertices. The mark \* indicates the results of this paper.

	natural		general	
	vertex guards	guards	vertex guards	guards
upper bound	does not exist [5]	$n - 2$ [*]	$n - 2$ [5]	$\lfloor (4n - 2)/5 \rfloor$ [*]
lower bound	does not exist [5]	$n - 2$ [*]	$\lfloor 2n/3 \rfloor - 1$ [3]	$\lceil (3n - 4)/5 \rceil$ [*]


## 2 Notation and Basic Properties

We are given a simple polygon  $P \subset \mathbb{R}^2$ . A *guard*  $g$  is a closed subset of the plane, whose boundary  $\partial g$  is described by a vertex  $v$  and two rays emanating from  $v$ . The ray that has the interior of the guard to its right is called the *left ray*, the other one is called the *right ray*. The *angle* of a guard is the interior angle formed by its rays. For a guard with angle  $\pi$ , the vertex is not unique.

A guard  $g$  *covers* an edge  $e$  of  $P$  *completely* if  $e \subseteq \partial g$  and their orientations match, that is, the inner side of  $e$  is on the inner side of  $g$ . We say  $e$  is covered *partly* by  $g$  if their orientations match and  $e \cap \partial g$  is a proper sub-segment of  $e$  that is not just a single point. We call a guard a  $k$ -guard if it covers exactly  $k$  edges completely. As  $P$  is simple, a guard can cover at most one edge partly. If a guard covers an edge partly and  $k$  edges completely, we call it a  $k'$ -guard. Assuming there are no collinear edges, a guard can cover at most two edges; then a natural vertex guard is a 2-guard and a natural edge guard is a 1-guard. A *guarding*  $\mathcal{G}(P)$  for  $P$  is a formula composed of a set of guards and the operators union and intersection that defines  $P$ . The *wireless localization problem* is to find a guarding with as few guards as possible. The same problem is sometimes referred to as *guard placement for point-in-polygon proofs* or the *sculpture garden problem* [5]. The following basic properties are restated without proof.

**Observation 1.** For any guarding  $\mathcal{G}(P)$  and for any two points  $p \in P$  and  $q \notin P$  there is a  $g \in \mathcal{G}(P)$  which distinguishes  $p$  and  $q$ , that is,  $p \in g$  and  $q \notin g$ .

**Lemma 1.** [4] Every edge of  $P$  must be covered by at least one guard or it must be covered partly by at least two guards.



**Fig. 1.** (a) a 2-guard, (b) a 1-guard (and a natural edge guard), (c) a 2-guard (and a natural vertex guard), (d) a 2-guard, (e) a 0-guard, (f) a 0-guard, (g) a 1-guard (not a 1'-guard), (h) a 1-guard (a non-natural vertex guard), (i) a 1'-guard.

### 3 Upper Bounds

Following Dobkin et al. [4] we use the notion of a *polygonal halfplane* which is a topological halfplane bounded by a *simple bi-infinite polygonal chain* with edges  $(e_1, \dots, e_n)$ , for  $n \in \mathbb{N}$ . For  $n = 1$ , the only edge  $e_1$  is a line and the polygonal halfplane is a halfplane. For  $n = 2$ ,  $e_1$  and  $e_2$  are rays which share a common source but are not collinear. For  $n \geq 3$ ,  $e_1$  and  $e_n$  are rays,  $e_i$  is a line segment, for  $1 < i < n$ , and  $e_i$  and  $e_j$ , for  $1 \leq i < j \leq n$ , do not intersect unless  $j = i + 1$  in which case they share an endpoint. For brevity we use the term *chain* in place of simple bi-infinite polygonal chain in the following. For a polygonal halfplane  $H$  define  $\gamma(H)$  to be the minimum integer  $k$  such that there exists a guarding  $\mathcal{G}(H)$  for  $H$  using  $k$  guards. Similarly, for a natural number  $n$ , denote by  $\gamma(n)$  the maximum number  $\gamma(H)$  over all polygonal halfplanes  $H$  that are bounded by a chain with  $n$  edges. Obviously  $\gamma(1) = \gamma(2) = 1$ . Dobkin et al. [4] show that  $\gamma(n) \leq n$ .

**Lemma 2.** *Any simple polygon  $P$  on  $n \geq 4$  vertices is an intersection of two polygonal halfplanes each of which consists of at least two edges.*

*Proof.* Let  $p_-$  and  $p_+$  be the vertices of  $P$  with minimal and maximal  $x$ -coordinate, respectively. If they are not adjacent along  $P$ , split the circular sequence of edges of  $P$  at both  $p_-$  and  $p_+$  to obtain two sequences of at least two segments each. Transform each sequence into a chain by linearly extending the first and the last segment beyond  $p_-$  or  $p_+$  to obtain a ray. As  $p_-$  and  $p_+$  are opposite extremal vertices of  $P$ , the two chains intersect exactly at these two points. Thus, the polygon  $P$  can be expressed as an intersection of two polygonal halfplanes bounded by these chains. Now consider the case that  $p_-$  and  $p_+$  are adjacent along  $P$ . Without loss of generality assume that  $P$  lies above the edge from  $p_-$  to  $p_+$ . Rotate clockwise until another point  $q$  has  $x$ -coordinate larger than  $p_+$ . If  $q$  and  $p_-$  are not adjacent along  $P$ , then split  $P$  at these points. Otherwise the convex hull of  $P$  is the triangle  $qp_-p_+$ . In particular,  $q$  and  $p_+$  are opposite non-adjacent extremal vertices and we can split as described above.  $\square$

**Theorem 3.** *Any simple polygon  $P$  with  $n \geq 4$  edges can be guarded using at most  $n - 2$  natural (vertex or edge) guards.*

*Proof.* Dobkin et al. [4] showed that for any chain there is a Peterson-style formula, that is, a guarding using natural edge guards only in which each guard appears exactly once and guards appear in the same order as the corresponding edges appear along the chain. Looking at the expression tree of this formula there is at least one vertex both of whose children are leaves. In other words, there is an operation (either union or intersection) that involves only two guards. As these two guards belong to two consecutive edges of  $P$ , we can replace this operation in the formula by the natural vertex guard of the common vertex, thereby saving one guard. Doing this for both chains as provided by Lemma 2 yields a guarding for  $P$  using  $n - 4$  natural edge guards and two natural vertex guards.  $\square$

The (closure of) the complement of a polygonal halfplane  $H$ , call it  $\overline{H}$ , is a polygonal halfplane as well.

**Observation 2.** *Any guarding for  $H$  can be transformed into a guarding for  $\overline{H}$  using the same number of guards.*

*Proof.* Use de Morgan's rules and invert all guards (keep their location but flip the angle to the complement with respect to  $2\pi$ ). Note that the resulting formula is monotone. Only guards complementary to the original ones appear (in SAT terminology: only negated literals); a formula is not monotone only if both a guard **and** its complementary guard appear in it.


**Corollary 4.** *Let  $P_1, \dots, P_m$  be a collection of  $m \geq 1$  simple polygons  $t$  of which are triangles, for  $0 \leq t \leq m$ . Let  $R$  be a region that can be described as a formula composed of the operations intersection, union, and complement over the variables  $\{P_1, \dots, P_m\}$  in which each  $P_i$  appears exactly once. Then  $R$  can be guarded using at most  $n - 2m + t$  natural (vertex or edge) guards, where  $n$  is the total number of edges of the polygons  $P_i$ , for  $1 \leq i \leq m$ .  $\square$*

**Corollary 5.** *Any simple polygon with  $n \geq 4$  edges and  $h$  non-triangular holes can be guarded using at most  $n - 2(h + 1)$  natural (vertex or edge) guards.  $\square$*

Our guarding scheme for chains is based on a recursive decomposition in which at each step the current chain is split into two or more subchains. At each split some segments are extended to rays and we have to carefully control the way these rays interact with the remaining chain(s). This is particularly easy if the split vertex lies on the convex hull because then the ray resulting from the segment extension cannot intersect the remainder of the chain at all. However, we have to be careful what we mean by convex hull. Instead of looking at the convex hull of a polygonal halfplane  $H$  we work with the convex hull of its bounding chain  $C$ . The convex hull  $h(C)$  of a chain  $C = (e_1, \dots, e_n)$ , for  $n \geq 2$ , is either the convex hull of  $H$  or the convex hull of  $\overline{H}$ , whichever of these two is not the whole plane which solely depends on the direction of the two rays of  $C$ . The boundary of  $h(C)$  is denoted by  $\partial h(C)$ . There is one degenerate case, when the two rays defining  $C$  are parallel and all vertices are contained in the strip between them; in this case,  $h(C)$  is a strip bounded by the two parallel lines through the rays and thus  $\partial h(C)$  is disconnected.

**Theorem 6.** *Any polygonal halfplane bounded by a simple bi-infinite polygonal chain with  $n \geq 2$  edges can be guarded using at most  $\lfloor (4n - 1)/5 \rfloor$  guards.*

*Proof.* The statement is easily checked for  $2 \leq n \leq 3$ . We proceed by induction on  $n$ . Let  $C$  be any chain with  $n \geq 4$  edges. Denote the sequence of edges along  $C$  by  $(e_1, \dots, e_n)$  and let  $v_i$ , for  $1 \leq i < n$ , denote the vertex of  $C$  incident to  $e_i$  and  $e_{i+1}$ . The underlying (oriented) line of  $e_i$ , for  $1 \leq i \leq n$ , is denoted by  $\ell_i$ . For  $2 \leq i \leq n - 1$ , let  $e_i^+$  be the ray obtained from  $e_i$  by extending the segment linearly beyond  $v_i$ . Similarly  $e_i^-$  refers to the ray obtained from  $e_i$  by extending the segment linearly beyond  $v_{i-1}$ . For convenience, let  $e_1^+ = \ell_1$  and  $e_n^- = \ell_n$ .




**Fig. 2.** The chain  $C$  can interact with the shaded region  $\Delta$  in three possible ways. The label  $\emptyset$  marks an area which does not contain any vertex from  $C$ .

Without loss of generality (cf. Observation 2) suppose that  $v_1$  is reflex, that is, the interior of the region bounded by  $C$  lies in the angle of  $C$  incident to  $v_1$  which is larger than  $\pi$ . If there is any vertex  $v_i$  on  $\partial h(C)$ , for some  $1 < i < n-1$ , then split  $C$  into two chains  $C_1 = (e_1, \dots, e_i^+)$  and  $C_2 = (e_{i+1}^-, \dots, e_n)$ . We obtain a guarding for  $C$  as  $\mathcal{G}(C_1) \cup \mathcal{G}(C_2)$  and thus  $\gamma(C) \leq \gamma(i) + \gamma(n-i)$ , for some  $2 \leq i \leq n-2$ . As both  $i \geq 2$  and  $n-i \geq 2$ , we can bound by the inductive hypothesis  $\gamma(C) \leq \lfloor (4i-1)/5 \rfloor + \lfloor (4n-4i-1)/5 \rfloor \leq \lfloor (4i-1)/5 + (4n-4i-1)/5 \rfloor \leq \lfloor (4n-1)/5 \rfloor$ . Else, if both  $e_1$  and  $e_n$  are part of  $\partial h(C)$  and  $\ell_1$  intersects  $\ell_n$  then we place a guard  $g$  that covers both rays at the intersection of  $\ell_1$  and  $\ell_n$  to obtain a guarding  $g \cup \mathcal{G}(e_2^-, \dots, e_{n-1}^+)$  for  $C$ . Therefore, in this case  $\gamma(C) \leq 1 + \gamma(n-2)$ . Observe that this is subsumed by the inequality from the first case with  $i = 2$ . Otherwise, either  $\ell_1$  does not intersect  $\ell_n$  and thus  $v_1$  and  $v_{n-1}$  are the only vertices of  $\partial h(C)$  (the degenerate case where  $\partial h(C)$  is disconnected) or without loss of generality (reflect  $C$  if necessary)  $v_1$  is the only vertex of  $\partial h(C)$ . Let  $\Delta$  denote the open wedge bounded by  $e_1$  and  $e_2^+$ . We distinguish three cases.


**Case 1.** There is a vertex of  $C$  in  $\Delta$  and among these, a vertex furthest from  $\ell_2$  is  $v_i$ , for some  $3 \leq i \leq n-2$  (Fig. 2(a)). Split  $C$  into three chains,  $C_1 = (\ell_1)$ ,  $C_2 = (e_2^-, \dots, e_i^+)$ , and  $C_3 = (e_{i+1}^-, \dots, e_n)$ . By the choice of  $v_i$  there is no intersection between  $C_2$  and  $C_3$  other than at  $v_i$ . A guarding for  $C$  can be obtained as  $\mathcal{G}(C_1) \cup (\mathcal{G}(C_2) \cap \mathcal{G}(C_3))$ . In this case  $\gamma(C) \leq 1 + \gamma(j) + \gamma(n-j-1)$ , for some  $2 \leq j \leq n-3$ . Since  $j \geq 2$  and  $n-j-1 \geq n-(n-3)-1 = 2$ , we can apply the inductive hypothesis to bound  $\gamma(C) \leq 1 + \lfloor (4j-1)/5 \rfloor + \lfloor (4n-4j-5)/5 \rfloor \leq \lfloor (4n-1)/5 \rfloor$ .

**Case 2.** There is a vertex of  $C$  in  $\Delta$  and among these, the unique one furthest from  $\ell_2$  is  $v_{n-1}$  (Fig. 2(b)). We may suppose that  $\ell_1$  intersects  $\ell_n$ ; otherwise (in the degenerate case where  $\partial h(C)$  is disconnected), exchange the roles of  $v_1$  and  $v_{n-1}$ . We cannot end up in Case 2 both ways. Let  $\Delta'$  denote the open (convex) wedge bounded by  $e_n$  and  $e_{n-1}^-$ . If there is any vertex of  $C$  in  $\Delta'$ , let  $v_i$  be such a vertex which is furthest from  $\ell_{n-1}$ . Let  $C_1 = (e_1, \dots, e_i^+)$  and  $C_2 = (e_{i+1}^-, \dots, e_{n-1}^+)$ .




Both  $C_1$  and  $C_2$  are simple, except that their first and their last ray may intersect (in that case split the resulting polygon into two chains). Put a guard  $g$  at the intersection of  $\ell_n$  with  $e_1$  such that  $g$  covers  $e_n$  completely and  $e_1$  partially (see figure, the small stripes indicate the side to be guarded). A guarding for  $C$  can be obtained as  $g \cap (\mathcal{G}(C_1) \cup \mathcal{G}(C_2))$ . Again this yields  $\gamma(C) \leq 1 + \gamma(i) + \gamma(n-i-1)$ , for some  $2 \leq i \leq n-3$ , and thus  $\gamma(C) \leq \lfloor (4n-1)/5 \rfloor$  as above in Case 1.

Otherwise there is no vertex of  $C$  in  $\Delta'$ . We distinguish two sub-cases. If  $e_{n-1}^+$  intersects  $e_1$  then put two guards (see figure): a first guard  $g_1$  at the intersection of  $\ell_n$  with  $e_1$  such that  $g_1$  covers  $e_n$  completely and  $e_1$  partially, and a second guard  $g_2$  at the intersection of  $\ell_{n-1}$  with  $e_1$  such that  $g_2$  covers  $e_{n-1}$  completely and  $e_1$  partially. Together  $g_1$  and  $g_2$  cover  $e_1$  and  $g_1 \cap (g_2 \cup \mathcal{G}(C'))$  provides a guarding for  $C$ , with  $C' = (e_2^-, \dots, e_{n-2}^+)$ . In this case we obtain  $\gamma(C) \leq 2 + \gamma(n-3)$  and thus by the inductive hypothesis  $\gamma(C) \leq 2 + \lfloor (4n-13)/5 \rfloor \leq \lfloor (4n-1)/5 \rfloor$ .




Finally, suppose that  $e_{n-1}^+$  does not intersect  $e_1$ . Then for the chain  $C' = (e_1, \dots, e_{n-1}^+)$  there is some vertex other than  $v_1$  on the convex hull boundary  $h(C')$ . Thus we can obtain a guarding for  $C'$  as described above for the case that there is more than one vertex on the convex hull. Put a guard  $g$  at the intersection of  $\ell_n$  with  $e_1$  such that  $g$  covers  $e_n$  completely and  $e_1$  partially (see figure). This yields a guarding  $g \cap \mathcal{G}(C')$  for  $C$  with  $\gamma(C) \leq 1 + \gamma(C') \leq 1 + \gamma(i) + \gamma(n-i-1)$ , for some  $2 \leq i \leq n-3$ . As in Case 1 we conclude that  $\gamma(C) \leq \lfloor (4n-1)/5 \rfloor$ .



**Case 3.** There is no vertex of  $C$  in  $\Delta$  (Fig. 2(c)). Let  $\Delta''$  denote the open (convex) wedge bounded by  $e_2^-$  and  $e_3^+$ . If  $e_3^-$  does not intersect  $e_1$  then put a natural vertex guard  $g$  at  $v_1$  to obtain a guarding  $g \cap \mathcal{G}(C')$  for  $C$ , where  $C' = (e_3^-, \dots, e_n)$ . This yields  $\gamma(C) \leq 1 + \gamma(n-2)$  and thus by the inductive hypothesis  $\gamma(C) \leq 1 + \lfloor (4n-9)/5 \rfloor \leq \lfloor (4n-1)/5 \rfloor$ .


Now suppose that  $e_3^-$  intersects  $e_1$ . We distinguish two sub-cases. If there is no vertex of  $C$  in  $\Delta''$ , then place two guards: a natural vertex guard  $g_1$  at  $v_1$  and a guard  $g_2$  at the intersection of  $e_3^-$  with  $e_1$  such that  $g_1$  covers  $e_3$  completely and  $e_1$  partially. A guarding for  $C$  is provided by  $g_1 \cap (g_2 \cup \mathcal{G}(C'))$ , with  $C' = (e_4^-, \dots, e_n)$ . In this case we obtain  $\gamma(C) \leq 2 + \gamma(n-3)$  and thus in the same way as shown above  $\gamma(C) \leq \lfloor (4n-1)/5 \rfloor$ .



Otherwise there is a vertex of  $C$  in  $\Delta''$ . Let  $v_i$ , for some  $4 \leq i \leq n-1$ , be a vertex of  $C$  in  $\Delta''$  which is furthest from  $\ell_3$ . First suppose  $e_{i+1}^-$  does not

intersect  $e_2$ . Then neither does  $e_i^+$  and hence we can split at  $v_i$  in the same way as if  $v_i$  would be on  $\partial h(C)$ . If  $i = n - 1$ ,  $e_n^-$  must intersect  $e_2$  (otherwise,  $e_n$  would be on  $\partial h(C)$ ). Thus we have  $i < n - 1$  and both chains consist of at least two segments/rays.

Now suppose that  $e_{i+1}^-$  intersects  $e_2$  and thus  $e_1$ , and denote the point of intersection between  $e_{i+1}^-$  and  $e_1$  by  $v'$ . Let  $e_1^*$  be the ray originating from  $v'$  in direction  $e_1$ , and let  $e_{i+1}^*$  denote the segment or ray (for  $i = n - 1$ ) originating from  $v'$  in direction  $e_{i+1}^-$ . Place a natural vertex guard  $g$  at  $v_1$ . Regardless of whether or not  $e_i^+$  intersects  $e_2$  and  $e_1$ , a guarding for  $C$  is provided by  $g \cap (\mathcal{G}(C_1) \cup \mathcal{G}(C_2))$ , with  $C_1 = (e_3^-, \dots, e_i^+)$  and  $C_2 = (e_1^*, e_{i+1}^*, \dots, e_n)$  (if  $i = n - 1$  then  $C_2 = (e_1^*, e_n^*)$ ). Observe that by the choice of  $v_i$  both  $C_1$  and  $C_2$  are simple and  $\gamma(C) \leq 1 + \gamma(j) + \gamma(n - j - 1)$ , for some  $2 \leq j \leq n - 3$ . As above, this yields  $\gamma(C) \leq \lfloor (4n - 1)/5 \rfloor$ .



We have shown that in every case  $\gamma(C) \leq \lfloor (4n - 1)/5 \rfloor$  and as  $C$  was arbitrary it follows that  $\gamma(n) \leq \lfloor (4n - 1)/5 \rfloor$ .  $\square$

**Corollary 7.** *Any simple polygon  $P$  with  $n$  edges can be guarded using at most  $\lfloor (4n - 2)/5 \rfloor$  guards.*

**Corollary 8.** *Let  $P_1, \dots, P_m$  be a collection of  $m \geq 1$  simple polygons with  $n$  edges in total, and let  $R$  be a region that can be described as a formula composed of the operations intersection, union, and complement over the variables  $\{P_1, \dots, P_m\}$  in which each  $P_i$  appears exactly once. Then  $R$  can be guarded using at most  $\lfloor (4n - 2m)/5 \rfloor$  guards.*

**Corollary 9.** *Let  $P$  be any simple polygon  $P$  with  $h$  holes such that  $P$  is bounded by  $n$  edges in total. Then  $P$  can be guarded using at most  $\lfloor (4n - 2h - 2)/5 \rfloor$  guards.*

## 4 Lower Bounds

For any natural number  $m$  we construct a polygon  $P_m$  with  $2m$  edges which requires “many” guards. The polygon consists of spikes  $S_1, S_2, \dots, S_m$  arranged in such a way that the lines through both edges of a spike cut into every spike to the left (see Fig. 3). Denote the apex of  $S_i$  by  $w_i$  and its left vertex by  $v_i$ . The edge from  $v_i$  to  $w_i$  is denoted by  $e_i$ , the edge from  $w_i$  to  $v_{i+1}$  by  $f_i$ . We can construct  $P_m$  as follows: Consider the two hyperbolas  $\{(x, y) \in \mathbb{R}^2 \mid x \geq 1, y = \frac{1}{x}\}$  and  $\{(x, y) \in \mathbb{R}^2 \mid x \geq 1, y = -\frac{1}{x}\}$ . Let  $v_1 := (1, 1)$  and  $w_1 := (1, -1)$ . Then choose  $f_1$  tangential to the lower hyperbola. Let  $v_2$  be the point where the tangent of the lower hyperbola intersects the upper hyperbola, that is,  $v_2 = (1 + \sqrt{2}, \frac{1}{1 + \sqrt{2}})$ . Choose  $w_2$  to be the point where the tangent of the upper hyperbola in  $v_2$  intersects the lower hyperbola, and proceed in this way. When reaching  $w_m$ , draw the last edge  $f_m$  from  $w_m$  to  $v_1$  to close the polygon. Due to the convexity of the hyperbolas,  $P_m$  has the claimed property.







Fig. 3. Example consisting of four spikes.

No two edges of  $P_m$  are collinear. Consider the line arrangement defined by the edges of  $P_m$ . No two lines intersect outside  $P_m$ , unless one of them is the line through  $f_m$ . This leads to the following observation.

**Observation 3.** *In any guarding for  $P_m$  every 2-guard that does not cover  $f_m$  is a natural vertex guard.*

**Theorem 10.** *For any even natural number  $n$  there exists a simple polygon with  $n$  edges which requires at least  $n - 2$  natural guards.*

We say a guard *belongs* to a spike  $S_i$  if it is a natural edge guard on  $e_i$  or  $f_i$  or if it is a natural vertex guard on  $v_i$  or  $w_i$ . As only natural guards are allowed, every guard belongs to exactly one spike. The basic idea is that most spikes must have at least two guards. Obviously every spike  $S_i$  has at least one guard, since  $e_i$  must be covered (Lemma 1).



**Lemma 11.** *Consider a guarding  $\mathcal{G}(P_m)$  using natural guards only, and let  $i \in \{1, \dots, m-1\}$ . If only one guard belongs to  $S_i$ , then this guard must be on  $v_i$  or on  $e_i$ . If neither the guard at  $w_i$  nor the guard of  $f_i$  appear in  $\mathcal{G}(P_m)$ , then both the guard at  $v_{i+1}$  and the guard of  $e_{i+1}$  are in  $\mathcal{G}(P_m)$ .*

*Proof.* Assume only one guard from  $\mathcal{G}(P_m)$  belongs to  $S_i$ . It cannot be the natural edge guard of  $f_i$ , because this would leave  $e_i$  uncovered (Lemma 1). If we had a guard on  $w_i$  only, there would be no guard to distinguish a point near  $v_i$  outside  $P_m$  from a point near  $v_{i+1}$  located inside  $P_m$  and below the line through  $f_i$  (see the two circles in the figure). Now assume there are no guards at  $w_i$  nor on  $f_i$ . Then to cover the edge  $f_i$  there must be a vertex guard on  $v_{i+1}$ . Furthermore, the edge guard on  $e_{i+1}$  is the only remaining natural guard to distinguish a point at the apex of  $S_i$  near  $w_i$  from a point located to the right of the apex of  $S_{i+1}$  near  $w_{i+1}$  and above the line through  $e_{i+1}$  (depicted by two crosses).  $\square$

This lemma immediately implies Theorem 10. Proceed through the spikes from left to right. As long as a spike has at least two guards which belong to it,

we are fine. Whenever there appears a spike  $S_i$  with only one guard, we know that there must be at least two guards in  $S_{i+1}$  namely at  $v_{i+1}$  and on  $e_{i+1}$ . Either there is a third guard that belongs to  $S_{i+1}$ , and thus both spikes together have at least four guards; or again we know already two guards in  $S_{i+2}$ . In this way, we can go on until we either find a spike which at least three guards belong to or we have gone through the whole polygon. So whenever there is a spike with only one guard either there is a spike with at least three guards that makes up for it, or every spike till the end has two guards. Hence there can be at most one spike guarded by one guard only that is not made up for later. For the last spike  $S_m$  the lemma does not hold and we only know that it has at least one guard. So all in all there are at least  $2(m-2) + 1 + 1 = n - 2$  guards.


If we allow general (vertex) guards, it is possible to find guardings for  $P_m$  using roughly  $2n/3$  guards, which is in accord with the lower bound in [3].

**Theorem 12.** *For any even natural number  $n$  there exists a simple polygon with  $n$  edges which requires at least  $\lceil (3n-4)/5 \rceil$  guards.*

*Proof.* Consider a polygon  $P_m$  as defined above, and let  $\mathcal{G}(P_m)$  be a guarding for  $P_m$ . Define  $a$  to be the number of 2-guards in  $\mathcal{G}(P_m)$ , and let  $b$  be the number of other guards. All the  $n$  edges of  $P$  have to be covered somehow. An edge can be covered completely by a 2-guard, a 1-guard, or a 1'-guard. If no guard covers it completely, then the edge must be covered by at least two guards partly (Lemma 1). Moreover, at least one of these guards, namely the one covering the section towards the right end of the edge, is a 0'-guard, because the orientation can not be correct to cover a second edge. So if an edge  $e$  is not covered by a 2-guard, then there is at least one guard that does not cover any edge other than  $e$ . Therefore  $2a + b \geq n$ .


For any  $i \in \{1, \dots, m-2\}$  let  $h_i$  be the directed line segment from the intersection of the lines through  $e_{i+1}$  and  $e_{i+2}$  to  $v_{i+2}$  (see Fig. 4). Similarly, let  $h'_i$  be the line segment from  $w_{i+1}$  to the intersection of the lines through  $f_i$  and  $f_{i+1}$ . As in Lemma 11, consider pairs  $(p_1, q_1), \dots, (p_{m-2}, q_{m-2})$  and  $(p'_1, q'_1), \dots, (p'_{m-2}, q'_{m-2})$  of points infinitesimally close to the starting point or the endpoint of the corresponding line segment, located as follows:  $p_i, p'_i \in P_m$  for all  $i$ ,  $q_i, q'_i \notin P_m$  for all  $i$ ,  $p_i$  is outside the natural vertex guard at  $w_{i+1}$ , whereas  $q_i$  is inside the natural vertex guard at  $w_{i+2}$ , and similarly,  $p'_i$  is outside the natural vertex guard at  $v_{i+2}$ , whereas  $q'_i$  is inside the natural vertex guard at  $v_{i+1}$ . There are  $n-4$  such pairs, and they need to be distinguished somehow (Observation 1). Any natural vertex guard can distinguish at most one pair, and the same is true for any (non-natural) 2-guard located along the line through  $f_m$ . Thus any 2-guard in  $\mathcal{G}(P_m)$  distinguishes at most one of the pairs (Observation 3).

We claim that every guard  $g$  in  $\mathcal{G}(P_m)$  can distinguish at most three of these pairs. Denote the vertex of  $g$  by  $v_g$ , and let  $\ell_g$  and  $r_g$  denote the left and right ray of  $g$ , respectively. Assume  $g$  distinguishes  $p_i$  from  $q_i$ . If  $v_g$  is to the left of  $h_i$ , then—in order to distinguish  $p_i$  from  $q_i$ —the ray  $r_g$  must intersect  $h_i$ . Symmetrically, if  $v_g$  is to the right of  $h_i$ , then  $\ell_g$  must intersect  $h_i$ . Finally, if  $v_g$  is on the line through  $h_i$  then it must be on the line segment  $h_i$  itself. To



**Fig. 4.** The pairs  $(p_i, q_i)$  and  $(p'_i, q'_i)$  must be distinguished.

distinguish  $p_i$  from  $q_i$ , the endpoint of  $h_i$  (i.e.  $v_{i+2}$ ) must be inside  $g$  (possibly on the boundary of  $g$ ), hence  $\ell_g$  must point to the left side of  $h_i$  or in the same direction as  $h_i$ , and  $r_g$  must point to the right side of  $h_i$  or in the same direction. Now assume  $g$  distinguishes  $p'_i$  and  $q'_i$ . If  $v_g$  is to the right of  $h'_i$ , then  $\ell_g$  must intersect it, if it is to the left  $r_g$  must intersect it. If  $v_g$  lies on  $h'_i$ ,  $\ell_g$  leaves to the left and  $r_g$  to the right, or either or both rays lie on  $h'_i$ . In any case either  $\ell_g$  intersects  $h_i$  ( $h'_i$ , respectively) coming from the right side of  $h_i$  ( $h'_i$ ) and leaving to the left side, or  $r_g$  intersects  $h_i$  ( $h'_i$ ) coming from the left side and leaving to the right, or  $\ell_g$  starts on  $h_i$  ( $h'_i$ ) itself leaving to the left or  $r_g$  starts on the line segment itself leaving to the right (see Fig. 5). If  $r_g$  leaves an oriented line segment to the right side of the segment or if  $\ell_g$  leaves an oriented line segment to the left side, we say the ray *crosses* the line segment *with correct orientation*. So whenever a pair  $(p_i, q_i)$  or  $(p'_i, q'_i)$  is distinguished by  $g$ , then at least one of the rays  $\ell_g$  or  $r_g$  has a correctly oriented crossing with  $h_i$  ( $h'_i$ , respectively). The line segments  $h_1, \dots, h_{m-2}$  lie on a oriented convex curve  $C$ , which we obtain by prolonging every line segment until reaching the starting point of the next one. Extend the first and last line segment to infinity vertically and horizontally, respectively. In the same way define a curve  $C'$  for  $h'_1, \dots, h'_{m-2}$  (see Fig. 4). Any ray can cross a convex curve at most twice. Because of the way  $C$  and  $C'$  are situated with respect to each other (a line that crosses  $C$  twice must have negative slope, to cross  $C'$  twice positive slope) a ray can intersect  $C \cup C'$  at most three times. But we are only interested in crossings with correct orientation. If a ray crosses a curve twice, exactly one of the crossings has the correct orientation. If a ray crosses both  $C$  and  $C'$  once, exactly one of the crossings has the correct orientation. Therefore any ray can have at most two correctly oriented crossings. If one of the rays has two correctly oriented crossings, the other ray has at most



**Fig. 5.** Different ways  $g$  can distinguish  $p$  and  $q$ . In every case  $\ell_g$  intersects  $h$  leaving to the left side or  $r_g$  intersects  $h$  leaving to the right.

one. Thus both rays together can have at most three correctly oriented crossings, and therefore  $g$  can distinguish at most three pairs. This leads to the second inequality  $a + 3b \geq n - 4$ . Both inequalities together imply  $a + b \geq \frac{3n-4}{5}$ .  $\square$

## References

1. Aigner, M., Ziegler, G.M.: Proofs from THE BOOK. Springer, Berlin, 3rd edition (2003)
2. Chvátal, V.: A Combinatorial Theorem in Plane Geometry. *J. Combin. Theory Ser. B* 18, 39–41 (1975)
3. Damian, M., Flatland, R., O’Rourke, J., Ramaswami, S.: A New Lower Bound on Guard Placement for Wireless Localization. 17th Annual Fall Workshop on Computational Geometry (2007)
4. Dobkin, D.P., Guibas, L., Hershberger, J., Snoeyink, J.: An Efficient Algorithm for Finding the CSG Representation of a Simple Polygon. *Algorithmica* 10,1–23 (1993)
5. Eppstein, D., Goodrich, M.T., Sitchinava, N.: Guard Placement for Efficient Point-in-Polygon Proofs. In: Proc. 23rd Annu. Sympos. Comput. Geom., pp. 27–36 (2007)
6. Fisk, S.: A Short Proof of Chvátal’s Watchman Theorem. *J. Combin. Theory Ser. B* 24, 374 (1978)
7. Fragoudakis, C., Markou, E., Zachos, S.: Maximizing the Guarded Boundary of an Art Gallery is APX-complete. *Comput. Geom. Theory Appl.* 38(3), 170–180 (2007)
8. Kahn, J., Klawe, M.M., Kleitman, D.J.: Traditional Galleries Require Fewer Watchmen. *SIAM J. Algebraic Discrete Methods* 4, 194–206 (1983)
9. Karavelas, M.I., Tsigaridas, E.P.: Guarding Curvilinear Art Galleries with Vertex or Point Guards. Rapport de recherche 6132, INRIA (2007)
10. Lee, D.T., Lin, A.K.: Computational Complexity of Art Gallery Problems. *IEEE Trans. Inform. Theory* 32(2), 276–282 (1986)
11. O’Rourke, J.: Galleries Need Fewer Mobile Guards: A Variation on Chvátal’s Theorem. *Geom. Dedicata* 14, 273–283, 1983.
12. O’Rourke, J.: Visibility. In: J. E. Goodman and J. O’Rourke, editors, *Handbook of Discrete and Computational Geometry*, chapter 28, pp. 643–663. CRC Press LLC, Boca Raton, FL (2004)
13. O’Rourke, J.: Computational Geometry Column 48. *ACM SIGACT News* 37(3), 55–57 (2006)
14. Shermer, T.C.: Recent Results in Art Galleries. *Proc. IEEE* 80(9), 1384–1399 (1992)
15. Speckmann, B., Tóth, C.D.: Allocating Vertex Pi-guards in Simple Polygons via Pseudo-triangulations. *Discrete Comput. Geom.* 33(2), 345–364 (2005)
16. Urrutia, J.: Art Gallery and Illumination Problems. In: J.-R. Sack and J. Urrutia, editors, *Handbook of Computational Geometry*, pp. 973–1027. Elsevier Science Publishers B.V. North-Holland, Amsterdam (2000)