Counting the Independent Sets of a Chordal Graph in Linear Time

Yoshio Okamoto	Toyohashi Univ Tech	JP
Takeaki Uno	National Inst Informatics	JP
Ryuhei Uehara	Japan Adv Inst Sci Tech	JP

June 6, 2005

4th Japanese-Hungarian Symposium on Discrete Math and Its Applications

Rényi Institute, Budapest, Hungary

An independent set

Setup $G = (V, E)$ a graph (undirected, finite, simple)

An independent set

Setup $G = (V, E)$ a graph (undirected, finite, simple)



An independent set

Setup

$G = (V, E)$ a graph (undirected, finite, simple)

Def.

A set $I \subseteq V$ is an **independent set** of G
if no two vertices in I are adjacent.



An independent set

Setup $G = (V, E)$ a graph (undirected, finite, simple)

Def. A set $I \subseteq V$ is an **independent set** of G if no two vertices in I are adjacent.



An independent set

Setup $G = (V, E)$ a graph (undirected, finite, simple)

Def. A set $I \subseteq V$ is an **independent set** of G if no two vertices in I are adjacent.



Not an independent set

An independent set

Setup $G = (V, E)$ a graph (undirected, finite, simple)

Def. A set $I \subseteq V$ is an **independent set** of G if no two vertices in I are adjacent.



Not an independent set

An independent set

Setup $G = (V, E)$ a graph (undirected, finite, simple)

Def. A set $I \subseteq V$ is an **independent set** of G if no two vertices in I are adjacent.



An independent set

Setup

$G = (V, E)$ a graph (undirected, finite, simple)

Def.

A set $I \subseteq V$ is an **independent set** of G if no two vertices in I are adjacent.



An independent set

An independent set

Setup $G = (V, E)$ a graph (undirected, finite, simple)

Def. A set $I \subseteq V$ is an **independent set** of G if no two vertices in I are adjacent.



An independent set

Also called a stable set of G

An enumeration problem

Input $G = (V, E)$ a graph

Output All independent sets of G



An enumeration problem

Input $G = (V, E)$ a graph

Output All independent sets of G



74 independent sets in total

A counting problem

Input $G = (V, E)$ a graph

Output # independent sets of G



A counting problem

Input	$G = (V, E)$ a graph
Output	# independent sets of G

Problem description

We study the following **counting problems**:

Input $G = (V, E)$ a graph

Output

(1) # independent sets of G

(2) # maximum independent sets of G

(3) # independent sets of G of fixed size

(4) # maximal independent sets of G

(5) # minimum maximal independent sets of G

A maximum independent set

Setup

$G = (V, E)$ a graph

Def.

An independent set I of G is **maximum**
if it has the largest size
among all independent sets of G.



A maximum independent set

Setup

$G = (V, E)$ a graph

Def.

An independent set I of G is **maximum**
if it has the largest size
among all independent sets of G.



Not maximum

A maximum independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **maximum**
if it has the largest size
among all independent sets of G.



A maximum independent set

Setup

$G = (V, E)$ a graph

Def.

An independent set I of G is **maximum**
if it has the largest size
among all independent sets of G.



Maximum

Maximum independent sets



74 independent sets

Maximum independent sets

74 independent sets

7 maximum independent sets

A maximal independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **maximal** if no proper superset of I is independent.



A maximal independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **maximal** if no proper superset of I is independent.



Maximal

A maximal independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **maximal** if no proper superset of I is independent.



A maximal independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **maximal** if no proper superset of I is independent.



Maximal

A maximal independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **maximal** if no proper superset of I is independent.



A maximal independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **maximal** if no proper superset of I is independent.



Not maximal

A maximal independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **maximal** if no proper superset of I is independent.



Not maximal

Maximal independent sets



74 independent sets

Maximal independent sets



74 independent sets

13 maximal independent sets

A minimum maximal independent set

Setup

$G = (V, E)$ a graph

Def.

An independent set I of G is **minimum maximal** if it is maximal and has the smallest size among all maximal independent sets of G.



maximal

A minimum maximal independent set

Setup

$G = (V, E)$ a graph

Def.

An independent set I of G is **minimum maximal** if it is maximal and has the smallest size among all maximal independent sets of G.



Not minimum maximal

A minimum maximal independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **minimum maximal** if it is maximal and has the smallest size among all maximal independent sets of G.



A minimum maximal independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **minimum maximal** if it is maximal and has the smallest size among all maximal independent sets of G.



Minimum maximal

A minimum maximal independent set

Setup $G = (V, E)$ a graph

Def. An independent set I of G is **minimum maximal** if it is maximal and has the smallest size among all maximal independent sets of G.



Minimum maximal

Also called a minimum independent dominating set of G

Minimum maximal independent sets



74 independent sets

Minimum maximal independent sets



74 independent sets

1 minimum maximal independent set

Problem description

We study the following **counting problems**:

Input $G = (V, E)$ a graph

Output

(1) # independent sets of G **74**

(2) # maximum independent sets of G **7**

(3) # independent sets of G of fixed size

(4) # maximal independent sets of G **13**

(5) # minimum maximal independent sets of G **1**

Independent sets of fixed size



74 independent sets

1 independent set of size 0

Independent sets of fixed size



74 independent sets

10 independent sets of size 1

Independent sets of fixed size



74 independent sets

29 independent sets of size 2

Independent sets of fixed size



74 independent sets

27 independent sets of size 3

Independent sets of fixed size

74 independent sets

7 independent sets of size 4

What's known

Fact These counting problems are #P-complete (analoguous to NP-completeness).

$\Longrightarrow$ Cannot hope for a poly-time algorithms.

What's known

Fact These counting problems are #P-complete (analoguous to NP-completeness).

$\Longrightarrow$ Cannot hope for a poly-time algorithms even for

- the line graphs of bipartite graphs (Valiant)
- bipartite graphs (Provan & Ball)
- planar bipartite graphs of max deg 4 (Vadhan).

What's known

Fact These counting problems are #P-complete (analoguous to NP-completeness).

$\Longrightarrow$ Cannot hope for a poly-time algorithms even for

- the line graphs of bipartite graphs (Valiant)
- bipartite graphs (Provan & Ball)
- planar bipartite graphs of max deg 4 (Vadhan).

$\Longrightarrow$ Focus on another class of perfect graphs!!

A chordal graph

Setup $G = (V, E)$ a graph

Def. G is **chordal**
if every induced cycle is of length three.

A chordal graph

Setup $G = (V, E)$ a graph

Def. G is **chordal**
if every induced cycle is of length three.



A chordal graph

Setup $G = (V, E)$ a graph

Def. G is **chordal**
if every induced cycle is of length three.



Not chordal

A chordal graph

Setup $G = (V, E)$ a graph

Def. G is **chordal**
if every induced cycle is of length three.



Not chordal

A chordal graph

Setup $G = (V, E)$ a graph

Def. G is **chordal**
if every induced cycle is of length three.



A chordal graph

Setup $G = (V, E)$ a graph

Def. G is **chordal**
if every induced cycle is of length three.



Chordal

Problem description

We study the following **counting problems**:

Input $G = (V, E)$ a chordal graph

Output

(1) # independent sets of G

(2) # maximum independent sets of G

(3) # independent sets of G of fixed size

(4) # maximal independent sets of G

(5) # minimum maximal independent sets of G

Results

We obtain the following results for chordal graphs.

(1) # independent sets of G
(2) # maximum independent sets of G
(3) # independent sets of G of fixed size

(4) # maximal independent sets of G
(5) # minimum maximal independent sets of G

Results

We obtain the following results for chordal graphs.

(1) # independent sets of G
(2) # maximum independent sets of G
(3) # independent sets of G of fixed size

$O(|V| + |E|)$ alg.

(4) # maximal independent sets of G
(5) # minimum maximal independent sets of G

#P-complete

Results

We obtain the following results for chordal graphs.

(1) # independent sets of G
(2) # maximum independent sets of G
(3) # independent sets of G of fixed size
$O(|V| + |E|)$ alg.
(4) # maximal independent sets of G
(5) # minimum maximal independent sets of G
#P-complete

Clique tree

Given

$G = (V, E)$ a (connected) chordal graph

Def

A **clique tree** of G is a tree T st

(1) the nodes of T = the maximal cliques of G,

(2) $\forall\, v \in V$,
the nodes of T containing v induce a tree.

Clique tree

Given $G = (V, E)$ a (connected) chordal graph

Def A **clique tree** of G is a tree T st

(1) the nodes of T = the maximal cliques of G,

(2) $\forall\, v \in V$,
the nodes of T containing v induce a tree.



Clique tree

Given $G = (V, E)$ a (connected) chordal graph

Def A **clique tree** of G is a tree T st

(1) the nodes of T = the maximal cliques of G,

(2) $\forall\, v \in V$,
the nodes of T containing v induce a tree.



Clique tree

Given $G = (V, E)$ a (connected) chordal graph

Def A **clique tree** of G is a tree T st

(1) the nodes of T = the maximal cliques of G,

(2) $\forall\, v \in V$,
the nodes of T containing v induce a tree.



Given $G = (V, E)$ a (connected) chordal graph

Def A **clique tree** of G is a tree T st

(1) the nodes of T = the maximal cliques of G,
(2) $\forall\, v \in V$,
the nodes of T containing v induce a tree.



Clique tree

Given $G = (V, E)$ a (connected) chordal graph

Def A **clique tree** of G is a tree T st

(1) the nodes of T = the maximal cliques of G,
(2) $\forall\, v \in V$,
the nodes of T containing v induce a tree.



A clique tree can be computed in linear time.

A basic property

Given $G = (V, E)$ a (connected) chordal graph,

A basic property

Given G = (V, E) a (connected) chordal graph,
T a clique tree of G, with root K

A basic property

Given $G = (V, E)$ a (connected) chordal graph,

T a clique tree of G, with root K





A basic property

Given $G = (V, E)$ a (connected) chordal graph,
T a clique tree of G, with root K

Property Every independent set of G either
contains exactly one vertex in K or not.





A basic property

Given $G = (V, E)$ a (connected) chordal graph,
T a clique tree of G, with root K

Property Every independent set of G either contains exactly one vertex in K or not.





Leads to a recursive formula...

Recursive formula

Let G be a chordal graph and T be a rooted clique tree of G. For a maximal clique K of G which is not a leaf of the clique tree, let $K_1, \ldots, K_\ell$ be the children of K in T. Furthermore, let $v \in K$. Then, the following identities hold.

$$\mathcal{IS}(G(K)) = \overline{\mathcal{IS}}(G(K), K) \dot\cup \dot\bigcup_{v \in K} \mathcal{IS}(G(K), v);$$

$$\mathcal{IS}(G(K), v) = \{S \cup \{v\} \mid S = \bigcup_{i=1}^{\ell} S_i, S_i \in \left\{ \begin{array}{ll} \mathcal{IS}(G(K_i), v) & \text{if } v \in K_i \\ \overline{\mathcal{IS}}(G(K_i), K \cap K_i) & \text{otherwise} \end{array} \right\}\};$$

$$\overline{\mathcal{IS}}(G(K), K) = \{S \mid S = \bigcup_{i=1}^{\ell} S_i, S_i \in \overline{\mathcal{IS}}(G(K_i), K \cap K_i)\};$$

$$\overline{\mathcal{IS}}(G(K_i), K \cap K_i) = \overline{\mathcal{IS}}(G(K_i), K_i) \dot\cup \dot\bigcup_{u \in K_i \setminus K} \mathcal{IS}(G(K_i), u) \quad \text{for each } i \in \{1, \ldots, \ell\},$$

where $\mathcal{IS}(G)$ denotes the family of independent sets in G, $\mathcal{IS}(G, v)$ denotes for a vertex v the family of independent sets in G including v, $\overline{\mathcal{IS}}(G, U)$ denotes the family of independent sets in G including no vertex of U for a vertex set U.

Recursive formula

Let G be a chordal graph and T be a rooted clique tree of G. For a maximal clique K of G which is not a leaf of the clique tree, let $K_1, \ldots, K_\ell$ be the children of K in T. Furthermore, let $v \in K$. Then, the following identities hold.

$$\mathcal{IS}(G(K)) = \overline{\mathcal{IS}}(G(K), K) \dot\cup \dot\bigcup_{v \in K} \mathcal{IS}(G(K), v);$$

$$\mathcal{IS}(G(K), v) = \{S \cup \{v\} \mid S = \bigcup_{i=1}^{\ell} S_i, S_i \in \left\{ \begin{array}{ll} \mathcal{IS}(G(K_i), v) & \text{if } v \in K_i \\ \overline{\mathcal{IS}}(G(K_i), K \cap K_i) & \text{otherwise} \end{array} \right\}\};$$

$$\overline{\mathcal{IS}}(G(K), K) = \{S \mid S = \bigcup_{i=1}^{\ell} S_i, S_i \in \overline{\mathcal{IS}}(G(K_i), K \cap K_i)\};$$

$$\overline{\mathcal{IS}}(G(K_i), K \cap K_i) = \overline{\mathcal{IS}}(G(K_i), K_i) \dot\cup \dot\bigcup_{u \in K_i \setminus K} \mathcal{IS}(G(K_i), u) \quad \text{for each } i \in \{1, \ldots, \ell\},$$

where $\mathcal{IS}(G)$ denotes the family of independent sets in G, $\mathcal{IS}(G, v)$ denotes for a vertex v the family of independent sets in G including v, $\overline{\mathcal{IS}}(G, U)$ denotes the family of independent sets in G including no vertex of U for a vertex set U.

$\Downarrow$

A detailed analysis yields a linear-time algorithm to count the independent sets in a chordal graph!

Recursive formula

Let G be a chordal graph and T be a rooted clique tree of G. For a maximal clique K of G which is not a leaf of the clique tree, let $K_1, \ldots, K_\ell$ be the children of K in T. Furthermore, let $v \in K$. Then, the following identities hold.

$$\mathcal{IS}(G(K)) = \overline{\mathcal{IS}}(G(K), K) \dot{\cup} \dot{\bigcup_{v \in K}} \mathcal{IS}(G(K), v);$$

$$\mathcal{IS}(G(K), v) = \{S \cup \{v\} \mid S = \bigcup_{i=1}^{\ell} S_i, S_i \in \left\{ \begin{array}{ll} \mathcal{IS}(G(K_i), v) & \text{if } v \in K_i \\ \overline{\mathcal{IS}}(G(K_i), K \cap K_i) & \text{otherwise} \end{array} \right\}\};$$

$$\overline{\mathcal{IS}}(G(K), K) = \{S \mid S = \bigcup_{i=1}^{\ell} S_i, S_i \in \overline{\mathcal{IS}}(G(K_i), K \cap K_i)\};$$

$$\overline{\mathcal{IS}}(G(K_i), K \cap K_i) = \overline{\mathcal{IS}}(G(K_i), K_i) \dot{\cup} \dot{\bigcup_{u \in K_i \setminus K}} \mathcal{IS}(G(K_i), u) \quad \text{for each } i \in \{1, \ldots, \ell\},$$

where $\mathcal{IS}(G)$ denotes the family of independent sets in G, $\mathcal{IS}(G, v)$ denotes for a vertex v the family of independent sets in G including v, $\overline{\mathcal{IS}}(G, U)$ denotes the family of independent sets in G including no vertex of U for a vertex set U.

$\Downarrow$

A detailed analysis yields a linear-time algorithm to count the independent sets in a chordal graph!

Thm The independent sets in a chordal graph $G = (V, E)$ can be counted in $O(|V| + |E|)$ time.

Plan

We obtain the following results for chordal graphs.

(1) # independent sets of G
(2) # maximum independent sets of G
(3) # independent sets of G of fixed size
$O(|V| + |E|)$ alg.
(4) # maximal independent sets of G
(5) # minimum maximal independent sets of G
#P-complete

Reduction of Set Cover

We use the following counting problem
(a counting version of the set cover problem).

Given	X a finite set, $\mathcal{S} \subseteq 2^X$ a family
Output	The number of subfamilies $\mathcal{S}' \subseteq \mathcal{S}$ st $\bigcup_{Y \in \mathcal{S}'} Y = X$.

Reduction of Set Cover

We use the following counting problem
(a counting version of the set cover problem).

Given X a finite set, $\mathcal{S} \subseteq 2^X$ a family

Output The number of subfamilies $\mathcal{S}' \subseteq \mathcal{S}$ st $\bigcup_{Y \in \mathcal{S}'} Y = X$.



Reduction of Set Cover

We use the following counting problem
(a counting version of the set cover problem).

Given X a finite set, $\mathcal{S} \subseteq 2^X$ a family

Output The number of subfamilies $\mathcal{S}' \subseteq \mathcal{S}$ st $\bigcup_{Y \in \mathcal{S}'} Y = X$.



Reduction of Set Cover

We use the following counting problem
(a counting version of the set cover problem).

Given X a finite set, $\mathcal{S} \subseteq 2^X$ a family

Output The number of subfamilies $\mathcal{S}' \subseteq \mathcal{S}$ st $\bigcup_{Y \in \mathcal{S}'} Y = X$.



Reduction of Set Cover

We use the following counting problem
(a counting version of the set cover problem).

Given X a finite set, $\mathcal{S} \subseteq 2^X$ a family

Output The number of subfamilies $\mathcal{S}' \subseteq \mathcal{S}$ st $\bigcup_{Y \in \mathcal{S}'} Y = X$.



Reduction of Set Cover

We use the following counting problem
(a counting version of the set cover problem).

Given X a finite set, $\mathcal{S} \subseteq 2^X$ a family

Output The number of subfamilies $\mathcal{S}' \subseteq \mathcal{S}$ st $\bigcup_{Y \in \mathcal{S}'} Y = X$.



Reduction of Set Cover

We use the following counting problem
(a counting version of the set cover problem).

Given X a finite set, $\mathcal{S} \subseteq 2^X$ a family

Output The number of subfamilies $\mathcal{S}' \subseteq \mathcal{S}$ st $\bigcup_{Y \in \mathcal{S}'} Y = X$.



Actual reduction



Actual reduction



Actual reduction



Actual reduction



Actual reduction



Actual reduction



$$\#\text{ set covers} + \sum_{x \in X} 2^{\#\text{ sets not containing } x}$$

$$\|$$

$$\#\text{ maximal independent sets}$$

Plan

We obtain the following results for chordal graphs.

(1) # independent sets of G
(2) # maximum independent sets of G
(3) # independent sets of G of fixed size
$O(|V| + |E|)$ alg.
(4) # maximal independent sets of G
(5) # minimum maximal independent sets of G
#P-complete

Plan

We obtain the following results for chordal graphs.

(1) # independent sets of G
(2) # maximum independent sets of G
(3) # independent sets of G of fixed size
$O(|V| + |E|)$ alg.
(4) # maximal independent sets of G
(5) # minimum maximal independent sets of G
#P-complete

[End of my talk]

Köszönöm szépen!